PREGABALIN capsules, for oral use, Initial U.S. Approval: 2004

Note: the content below do not include all the information needed to use Pregabalin capsules safely and effectively. Please read the package insert carefully and follow the doctors' instructions before use the product.

	INDICATIONS	AND	USAGE	
Progabalia Canculas is indicate	nd for			

- Pregabalin Capsules is indicated for:
- Neuropathic pain associated with diabetic peripheral neuropathy (DPN)
- Postherpetic neuralgia (PHN)
- Adjunctive therapy for adult patients with partial onset seizures
- Fibromyalgia
- Neuropathic pain associated with spinal cord injury

------ DOSAGE AND ADMINISTRATION ------

- For all indications, begin dosing at 150 mg/day.
- Dosing recommendations:

INDICATION	Dosing Regimen	Maximum Dose
DPN Pain (2.1)	3 divided doses per day	300 mg/day within 1 week
PHN (2.2)	2 or 3 divided doses per day	300 mg/day within 1 week. Maximum dose of 600 mg/day.
Adjunctive Therapy for Adult Patients with Partial Onset Seizures (2.3)	2 or 3 divided doses per day	Maximum dose of 600 mg/day.
Fibromyalgia (2.4)	2 divided doses per day	300 mg/day within 1 week. Maximum dose of 450 mg/day.
Neuropathic Pain Associated with Spinal Cord Injury (2.5)	2 divided doses per day	300 mg/day within 1 week. Maximum dose of 600 mg/day.

Dose should be adjusted in patients with reduced renal function.
DOSAGE FORMS AND STRENGTHS
Capsules: 75 mg and 300 mg.